

Exploring Born-Infeld electrodynamics using plasmas

DA Burton*

RMGM Trines[†]

TJ Walton*

H Wen*

June 14, 2010

Abstract

The behaviour of large amplitude electrostatic waves in cold plasma is investigated in the context of Born-Infeld electrodynamics. The equations of motion for the plasma are established using an unconstrained action principle. The maximum amplitude and frequency of a large amplitude electrostatic wave are determined, and a lower bound on the wavelength is established. The maximum electric field is found to be the same as that on a point electron at rest.

1 Introduction

Recent years have seen renewed interest in non-linear electrodynamics [1–3], in particular Born-Infeld theory [4], and its implications. Unlike their Maxwell counterparts, the Born-Infeld field equations are *fundamentally* non-linear; they are non-linear even in the vacuum. Born and Infeld introduced their theory [4] in order to ameliorate the singular self-energy of a point charge. Furthermore, it was discovered that among the family of non-linear generalizations of Maxwell electrodynamics, Born-Infeld theory possesses a number of highly attractive features; in particular, like the vacuum Maxwell equations, the vacuum Born-Infeld equations exhibit zero birefringence and its solutions have exceptional causal behaviour [5, 6]. Furthermore, Born-Infeld theory shares a number of properties with the low energy dynamics of strings and branes [7].

Some of the most extreme conditions ever encountered in a terrestrial laboratory are created when high-power laser pulses interact with matter. The laser pulse immediately vaporizes the matter to form an intense laser-plasma providing novel avenues for generating intense bursts of coherent electromagnetic radiation for a wide range of applications in biological and material science [8]. Furthermore, laser-plasmas permit controllable investigation of matter in extreme conditions that only occur naturally away from the Earth. It is expected that the next generation of ultra-intense lasers will, for the first time, allow controllable access to regimes where a host of different quantum electrodynamic phenomena will be evident [9] and theories of non-linear electrodynamics will be central to their study. However, it is conceivable that Born-Infeld electrodynamics may play a role before it is necessary to invoke quantum electrodynamics [3].

Motivated by recent analyses of the propagation of Born-Infeld electromagnetic waves in waveguides [2] and in background uniform magnetic fields [3], this article addresses aspects of the non-linear electrodynamics of laser-plasmas.

A sufficiently short and intense laser pulse propagating through a plasma may create a travelling longitudinal plasma wave whose velocity is approximately the same as the laser pulse's group velocity. However, it is not possible to sustain arbitrarily large electric fields; substantial numbers of plasma electrons become trapped in the wave and are accelerated, which dampens the wave (the wave 'breaks'). Early theoretical investigation of non-linear plasma waves was undertaken by Akhiezer and Polovin [10], and later expounded by Dawson [11] in the context of wave-breaking.

Wave-breaking is a fundamentally non-linear phenomenon, and it is natural to explore the properties of Born-Infeld plasmas from this perspective. For simplicity, in the present article we adopt the *cold* plasma model for the electron fluid and generalize well-known results [10, 11], for the cold Maxwell plasma, to the context

*Department of Physics, Lancaster University, UK and the Cockcroft Institute, UK

[†]Rutherford Appleton Laboratory, Chilton, Didcot, UK and Department of Physics, Lancaster University, UK.

of non-linear electrodynamics. In particular, we obtain an exact expression for the maximum electric field of electrostatic oscillations in a cold Born-Infeld plasma and analyse the wavelength and frequency of those oscillations. There are numerous discussions of electrostatic waves in a warm Maxwell plasma; for example, see [12, 13].

We employ the Einstein summation convention throughout this article. Latin indices a, b, c run over $0, 1, 2, 3$ and units are used in which the speed of light $c = 1$ and the permittivity of the vacuum $\varepsilon_0 = 1$.

Let (x^a) be an inertial coordinate system on Minkowski spacetime $(\mathcal{M}, g)$ where x^0 is the proper time of observers at fixed Cartesian coordinates (x^1, x^2, x^3) in the laboratory. The metric tensor g has the form

$$g = \eta_{ab} dx^a \otimes dx^b \quad (1)$$

with

$$\eta_{ab} = \begin{cases} -1 & \text{if } a = b = 0, \\ 1 & \text{if } a = b \neq 0, \\ 0 & \text{if } a \neq b \end{cases} \quad (2)$$

and the Hodge map $\star$ is induced from the 4-form $\star 1$ on $\mathcal{M}$ where

$$\star 1 = dx^0 \wedge dx^1 \wedge dx^2 \wedge dx^3. \quad (3)$$

The plasma electrons are represented as a cold relativistic fluid; their worldlines are trajectories of the unit normalized future-pointing timelike 4-vector field V on $\mathcal{M}$ and the 0-form n is their proper number density. We are interested in the evolution of a plasma over timescales during which the motion of the ions is negligible in comparison with the motion of the electrons, and we assume that the ions are at rest and distributed homogeneously in the laboratory frame. Their worldlines are trajectories of the vector field $N_{\text{ion}} = n_{\text{ion}} \partial / \partial x^0$ on $\mathcal{M}$ where n_{ion} is the ion number density (a positive constant) in the laboratory frame.

Maxwell's equations may be written covariantly as

$$dF = 0, \quad d \star G = -qn \star \tilde{V} - q_{\text{ion}} \star \widetilde{N_{\text{ion}}} \quad (4)$$

where $q < 0$ is the charge on the electron and $q_{\text{ion}} = Z|q|$ is the charge on an ion with Z the multiplicity of the ionization. The 1-forms $\tilde{V}$ and $\widetilde{N_{\text{ion}}}$ are the metric duals of the vector fields V and N_{ion} respectively, i.e. the 1-form $\tilde{V}$ satisfies $\tilde{V}(U) = g(V, U)$ for all vector fields U on $\mathcal{M}$. The Maxwell 2-form F encodes the electric field $\mathbf{E}$ and the magnetic induction $\mathbf{B}$; the excitation 2-form G encodes the electric displacement $\mathbf{D}$ and the magnetic field $\mathbf{H}$. In Maxwell electrodynamics $G = F$ in the vacuum; however, in Born-Infeld electrodynamics (4) is retained but the constitutive relations for $(\mathbf{D}, \mathbf{H})$ in terms of $(\mathbf{E}, \mathbf{B})$ are non-linear in the vacuum. In Born-Infeld electrodynamics the constitutive relation for G in terms of F is

$$G = \frac{1}{\sqrt{1 - \kappa^2 X - \kappa^4 Y^2/4}} (F - \kappa^2 Y \star F) \quad (5)$$

where the invariants X and Y are

$$X = \star(F \wedge \star F), \quad Y = \star(F \wedge F) \quad (6)$$

and κ is a new constant of nature.

To motivate the field equations used herein, we begin section 2 by developing them using an unconstrained action principle [14]. We will briefly review how (5) arises and show that V satisfies the field equations

$$\nabla_V \tilde{V} = \frac{q}{m} \iota_V F, \quad g(V, V) = -1 \quad (7)$$

for a relativistic cold electron fluid, where $q\iota_V F$ is the Lorentz 4-force acting on the electron fluid, ι_V is the interior product with respect to V , m is the electron rest mass and ∇ is the Levi-Civita connection on $\mathcal{M}$. This will be followed in section 3 by an exploration of the properties of large amplitude relativistic electrostatic oscillations.

2 Action principle

A succinct action principle for establishing (4, 7) is formulated using a *three*-dimensional manifold $\mathcal{B}$ (often called a ‘material’ [15], or ‘body’, manifold), where each point in $\mathcal{B}$ corresponds to an integral curve of V .

The independent variables of the action introduced below are the electromagnetic 1-form A on $\mathcal{M}$ and a submersion f from $\mathcal{M}$ to $\mathcal{B}$,

$$f : \mathcal{M} \longrightarrow \mathcal{B} \quad (8)$$

$$x^a \longmapsto \xi^A = f^A(x). \quad (9)$$

The map f identifies an integral curve of V with its corresponding point in $\mathcal{B}$, where upper case Latin indices $A, B = 1, 2, 3$. The inverse image $f^{-1}(P)$ of the point $P \in \mathcal{B}$ is the worldline of an idealized ‘material’ particle in the electron fluid and V may be determined from f as the unique timelike future-pointing vector field satisfying

$$df^A(V) = 0 \quad (10)$$

and

$$g(V, V) = -1. \quad (11)$$

In the present approach, material properties of the electron fluid are encoded in tensors and forms on $\mathcal{B}$, and fields on $\mathcal{M}$ are induced from material tensors on $\mathcal{B}$ using f . In particular, the number of electrons $N[\Sigma]$ occupying the spacelike hypersurface Σ may be induced from a 3-form Ω on $\mathcal{B}$ as

$$N[\Sigma] = \int_{\Sigma} f^* \Omega \quad (12)$$

where f^* denotes the pull-back map, induced from f , on forms on $\mathcal{B}$ to forms on $\mathcal{U}$. The electron number current 3-form j ,

$$j = f^* \Omega \quad (13)$$

is identically closed,

$$dj = 0, \quad (14)$$

where d is the exterior derivative, which follows because Ω is a top form on $\mathcal{B}$ and $df^* = f^*d$.

A variational principle yielding (7) for V and (4) for the potential 1-form A , where $F = dA$, employs the action functional S where

$$S[A, f] = \int_{\mathcal{M}} [\mathcal{L}_{\text{EM}}(X, Y) \star 1 - m\sqrt{j \cdot j} \star 1 - qA \wedge j - q_{\text{ion}}A \wedge j_{\text{ion}}]. \quad (15)$$

The 0-form $\mathcal{L}_{\text{EM}}(X, Y)$ is a local function of the invariants X and Y only. For standard Maxwell electrodynamics $\mathcal{L}_{\text{EM}} = X/2$, whereas for Born-Infeld electrodynamics $\mathcal{L}_{\text{EM}} = \mathcal{L}_{\text{BI}}$ and

$$\mathcal{L}_{\text{BI}}(X, Y) = \frac{1}{\kappa^2} (1 - \sqrt{1 - \kappa^2 X - \kappa^4 Y^2/4}). \quad (16)$$

The 3-form $j_{\text{ion}} = n_{\text{ion}} dx^1 \wedge dx^2 \wedge dx^3$ is the background ion number current and the 0-form $j \cdot j$ is the square of the magnitude of j ,

$$j \cdot j = \star^{-1}(j \wedge \star j) \quad (17)$$

where $\star^{-1}$ is the inverse of the Hodge map $\star$.

The first term $\mathcal{L}_{\text{EM}}(X, Y) \star 1$ in the Lagrangian 4-form in (15) depends only on A and g , the second term $m\sqrt{j \cdot j} \star 1$ depends only on f and g , the third term $qA \wedge j$ couples f and A and the fourth term $q_{\text{ion}}A \wedge j_{\text{ion}}$ depends only on A .

We will show in the following that the field equations (4, 7) for n , V and A are recovered from those for f and A by introducing the electron proper number density n and 4-velocity field V as

$$n = \sqrt{j \cdot j}, \quad (18)$$

$$\tilde{V} = \frac{\star^{-1}j}{n}. \quad (19)$$

2.1 Non-linear generalization of the Maxwell equations

Equations for the electromagnetic field F arise upon seeking stationary variations of S with respect to A :

$$\delta_A S = \int_{\mathcal{M}} \left[\left(\frac{\partial \mathcal{L}_{\text{EM}}}{\partial X} \delta_A X + \frac{\partial \mathcal{L}_{\text{EM}}}{\partial Y} \delta_A Y \right) \star 1 - \delta A \wedge (qj + q_{\text{ion}} j_{\text{ion}}) \right] \quad (20)$$

where $\delta_A \alpha$ denotes the variation of a form α with respect to A . Using (6) and $\star \star \alpha = -\alpha$ where α is a 4-form on $\mathcal{M}$, it follows

$$X \star 1 = -F \wedge \star F, \quad Y \star 1 = -F \wedge F \quad (21)$$

and

$$\delta_A X \star 1 = \delta_A (X \star 1) = -2d\delta A \wedge \star F, \quad \delta_A Y \star 1 = \delta_A (Y \star 1) = -2d\delta A \wedge F. \quad (22)$$

Hence

$$\delta_A S = \int_{\mathcal{M}} \delta A \wedge \left[-2d \left(\frac{\partial \mathcal{L}_{\text{EM}}}{\partial X} \star F + \frac{\partial \mathcal{L}_{\text{EM}}}{\partial Y} F \right) - (qj + q_{\text{ion}} j_{\text{ion}}) \right] \quad (23)$$

where the variation δA is chosen to have compact support on $\mathcal{M}$, and an integration by parts has been used in the final step. By demanding that the action is stationary under all such variations we obtain

$$d \star G = -qj - q_{\text{ion}} j_{\text{ion}} \quad (24)$$

where

$$\star G = 2 \left(\frac{\partial \mathcal{L}_{\text{EM}}}{\partial X} \star F + \frac{\partial \mathcal{L}_{\text{EM}}}{\partial Y} F \right). \quad (25)$$

The field equations (4) are obtained by introducing n and V using (18) and (19), and the ion number 4-current N_{ion} as

$$\widetilde{N_{\text{ion}}} = \star^{-1} j_{\text{ion}} \quad (26)$$

and noting $dF = 0$ since F is an exact 2-form. Equation (5) immediately follows from (16, 25) with $\mathcal{L}_{\text{EM}} = \mathcal{L}_{\text{BI}}$ and using $\star \star \beta = -\beta$ where β is a 2-form.

2.2 Field equations for the electron fluid

The field equations (7) describing the electron fluid are obtained by seeking stationary variations of S with respect to f . Using (18) it follows

$$\delta_f \sqrt{j \cdot j} = \frac{1}{\sqrt{j \cdot j}} [\star^{-1} (\delta_f j \wedge \star j)]. \quad (27)$$

To proceed further we express $j = f^* \Omega$ explicitly in terms of the components $\{f^A\}$ of f .

$$\begin{aligned} \delta_f j &= \frac{1}{3!} \left(\frac{\partial \Omega_{ABC}}{\partial \xi^D} \circ f \right) \delta f^D df^A \wedge df^B \wedge df^C + \frac{1}{2!} (\Omega_{ABC} \circ f) d\delta f^A \wedge df^B \wedge df^C \\ &= \iota_W dj + d\iota_W j \end{aligned} \quad (28)$$

where $\Omega = \frac{1}{3!}\Omega_{ABC}(\xi) d\xi^A \wedge d\xi^B \wedge d\xi^C$ and

$$j = f^*\Omega = \frac{1}{3!}(\Omega_{ABC} \circ f) df^A \wedge df^B \wedge df^C \quad (29)$$

have been used. The vector field W is

$$W = \delta f^A W_A \quad (30)$$

where $\{W_A\}$ is a basis for V -orthogonal vector fields on $\mathcal{M}$, and the frame $\{V, W_A\}$ and coframe $\{-\tilde{V}, df^A\}$ are naturally dual, i.e.

$$df^A(W_B) = \delta_B^A, \quad \tilde{V}(W_A) = 0, \quad df^A(V) = 0, \quad \tilde{V}(V) = -1 \quad (31)$$

where δ_B^A is the Kronecker delta. Since j is a closed 3-form it follows

$$\delta_f j = d\iota_W j \quad (32)$$

using (28) and so

$$\begin{aligned} \delta_f(\sqrt{j \cdot j} \star 1) &= \frac{1}{\sqrt{j \cdot j}} \delta_f j \wedge \star j \\ &= d\iota_W j \wedge \frac{1}{\sqrt{j \cdot j}} \star j. \end{aligned} \quad (33)$$

Varying (15) with respect to f yields

$$\delta_f S = - \int_{\mathcal{M}} \left[\frac{m}{\sqrt{j \cdot j}} \delta_f j \wedge \star j + qA \wedge \delta_f j \right] \quad (34)$$

where the variation δf^A is chosen to have compact support on $\mathcal{M}$. An integration by parts yields

$$\delta_f S = \int_{\mathcal{M}} \iota_W j \wedge d\left(\frac{m}{\sqrt{j \cdot j}} \star j - qA\right) \quad (35)$$

and by demanding that S is stationary under all suitable variations δf^A we obtain

$$\iota_{W_A} j \wedge d\left(\frac{m}{\sqrt{j \cdot j}} \star j - qA\right) = 0. \quad (36)$$

Eliminating j in favour of n and V using (18, 19) yields

$$\iota_{W_A} \iota_V (m d\tilde{V} - qF) = 0 \quad (37)$$

and since $\{V, W_A\}$ is a frame on $\mathcal{M}$ it follows

$$\iota_V d\tilde{V} = \frac{q}{m} \iota_V F. \quad (38)$$

Since $g(V, V)$ is constant it may be shown $\iota_V d\tilde{V} = \nabla_V \tilde{V}$ (see, for example, [16]) and (7) follows immediately.

3 Non-linear electrostatic oscillations

The remainder of this article focuses on properties of large-amplitude longitudinal electrostatic waves propagating parallel to the x^3 -axis with phase velocity v (with $0 < v < 1$) in the laboratory frame. We introduce the pair $\{e^1, e^2\}$

$$e^1 = v dx^3 - dx^0, \quad e^2 = dx^3 - v dx^0 \quad (39)$$

and note that the orthonormal coframe $\{\gamma e^1, \gamma e^2, dx^1, dx^2\}$ is adapted to observers moving at velocity v along x^3 (i.e observers in the ‘wave frame’). We seek a 4-velocity V of the form

$$\tilde{V} = \mu(\zeta)e^1 + \psi(\zeta)e^2 \quad (40)$$

where $\zeta = x^3 - vx^0$ is the wave’s phase and we have adopted the so-called ‘quasi-static approximation’; the pointwise dependence of μ and ψ is on ζ only. Using $g(V, V) = -1$ the component ψ is found as

$$\psi = -\sqrt{\mu^2 - \gamma^2} \quad (41)$$

where the Lorentz factor $\gamma = 1/\sqrt{1-v^2}$ and the sign of ψ is chosen to ensure that the velocity $\gamma e^2(V)$ of the electron fluid in the wave frame is non-positive. Thus, in the laboratory frame the speed of the electrons is slower than the phase speed of the wave, except at wave-breaking where the electrons catch the wave.

In the present analysis we assume that the electromagnetic field is due entirely to the electron fluid and ion background. The magnetic field vanishes and the only non-zero component of the electric field is in the x^3 direction; it follows that the Maxwell 2-form F is

$$F = E dx^0 \wedge dx^3 \quad (42)$$

where E is the x^3 component of the electric field.

Since $\tilde{V}$ and F are elements of the subspace of forms on $\mathcal{M}$ generated by $\{dx^0, dx^3\}$ it follows

$$\tilde{V} \wedge (d\tilde{V} - \frac{q}{m}F) = 0 \quad (43)$$

and, using (38, 43) with $g(V, V) = -1$, we obtain

$$d\tilde{V} = \frac{q}{m}F \quad (44)$$

and (40, 42, 44) yield

$$E = \frac{1}{\gamma^2} \frac{m}{q} \frac{d\mu}{d\zeta}. \quad (45)$$

Since $d\zeta \wedge F = 0$ and $dF = 0$ it follows

$$d\left(\frac{\partial \mathcal{L}_{\text{EM}}}{\partial Y} F\right) = 0 \quad (46)$$

and inserting (45, 40) into (4) yields

$$\frac{d}{d\zeta} \left(2 \frac{\partial \mathcal{L}_{\text{EM}}}{\partial X} \Big|_{Y=0} \frac{d\mu}{d\zeta} \right) = \frac{q^2}{m} (n\mu - Zn_{\text{ion}}), \quad n = \frac{Zn_{\text{ion}}\gamma^2 v}{\sqrt{\mu^2 - \gamma^2}} \quad (47)$$

where

$$Y = 0 \quad (48)$$

follows from (6) and (46) has been used. Hence

$$\frac{d}{d\zeta} \left(2 \frac{\partial \mathcal{L}_{\text{EM}}}{\partial X} \Big|_{Y=0} \frac{d\mu}{d\zeta} \right) = \frac{q^2}{m} Zn_{\text{ion}} \gamma^4 \left(\frac{v\mu}{\sqrt{\mu^2 - \gamma^2}} - 1 \right). \quad (49)$$

The product of (49) and $d\mu/d\zeta$ may be written

$$\frac{d}{d\zeta} \left[\left(2 \frac{\partial \mathcal{L}_{\text{EM}}}{\partial X} X - \mathcal{L}_{\text{EM}} \right) \Big|_{Y=0} - mZn_{\text{ion}}(v\sqrt{\mu^2 - \gamma^2} - \mu) \right] = 0 \quad (50)$$

using

$$X = E^2, \quad (51)$$

which follows from (6).




Figure 1: The dashed curve shows μ versus ζ and the solid curve shows E versus ζ (not to scale). Points of intersection of E with the ζ -axis are labeled I and III, and II is a turning point of E . Using (45) and $q < 0$ it follows $d\mu/d\zeta$ and E are of opposite sign.

3.1 Maximum amplitude oscillation

The square root in the right-hand side of (49) places a lower bound on μ . Suppose that over an oscillation the component μ of $\tilde{V}$ attains its lowest possible value $\mu_I = \gamma$ at $\zeta = \zeta_I$ (see figure 1). Since μ_I is a turning point of μ it follows $E(\zeta_I) = 0$. At $\zeta = \zeta_{II}$ the electric field $E(\zeta_{II}) = -E_{\max}$ where $E_{\max}$ is the amplitude of the largest possible oscillation (see figure 1). Integration of (50) over $[\zeta_I, \zeta_{II}]$ yields

$$\left(2\frac{\partial \mathcal{L}_{\text{EM}}}{\partial X}X - \mathcal{L}_{\text{EM}}\right)\bigg|_{\substack{X=E_{\max}^2 \\ Y=0}} + \mathcal{L}_{\text{EM}}\bigg|_{\substack{X=0 \\ Y=0}} = mZn_{\text{ion}}(\gamma - 1) \quad (52)$$

For a standard cold Maxwell plasma $\mathcal{L}_{\text{EM}} = X/2$, which inserted into (52) yields the well-known wave-breaking limit due to Akhiezer and Polovin [10]

$$E_{\max}^{\text{AP}} = \sqrt{2mZn_{\text{ion}}(\gamma - 1)} \quad (53)$$

$$= \frac{m\omega_p c}{|q|} \sqrt{2(\gamma - 1)} \quad (54)$$

for a multiply-ionized relativistic cold plasma, where

$$\omega_p = \sqrt{\frac{q^2 Zn_{\text{ion}}}{m\varepsilon_0}} \quad (55)$$

is the plasma frequency and the speed of light c and permittivity of the vacuum ε_0 have been restored.

Using $\mathcal{L}_{\text{EM}} = \mathcal{L}_{\text{BI}}$ and (16, 52, 53) it follows

$$E_{\max}^{\text{BI}} = \frac{1}{\kappa} \sqrt{1 - [\kappa^2 (E_{\max}^{\text{AP}})^2 / 2 + 1]^{-2}} \quad (56)$$

for a cold Born-Infeld plasma and $\lim_{\kappa \rightarrow 0} E_{\max}^{\text{BI}} = E_{\max}^{\text{AP}}$ as expected.

The choice $\kappa = \varepsilon_0 r_0^2 / |q|$ made by Born and Infeld, where r_0 is the classical radius of the electron, leads to $\kappa \sim 10^{-22} \text{ m/V}$. However, quantum electrodynamic vacuum effects are expected to become important

at electric field strengths $\sim 10^{18}$ V/m, and if Born-Infeld theory plays a role at the classical level it follows $E_{\max}^{\text{BI}} \lesssim 10^{18}$ V/m and $\kappa E_{\max}^{\text{BI}} \ll 1$. The next generation of laser systems (such as Extreme Light Infrastructure (ELI) [17] and High Power Laser Energy Research system (HiPER) [18]), are expected to offer intensities $\sim 10^{25}$ W/cm² corresponding to electric field strengths $\sim 10^{16}$ V/m.

Although in practice $\kappa E_{\max}^{\text{BI}} \ll 1$ it is worth noting that although the limit of $E_{\max}^{\text{AP}}$ as $v \rightarrow c$ does not exist, using (56) it follows

$$\lim_{v \rightarrow c} E_{\max}^{\text{BI}} = \frac{1}{\kappa} \quad (57)$$

which is the value of the electric field of a static point electron evaluated at the location of the point electron in Born-Infeld electrodynamics [4].

3.2 Period and frequency of maximum amplitude oscillations

The period λ of the maximum amplitude oscillation of a cold Born-Infeld plasma is obtained from the solution to (49) with $\mathcal{L}_{\text{EM}} = \mathcal{L}_{\text{BI}}$ and the initial conditions $\mu(\zeta_{\text{I}}) = \gamma$ and $d\mu/d\zeta|_{\zeta=\zeta_{\text{I}}} = 0$. The particular first integral of (49) satisfying the initial conditions on μ may be written

$$\left(\frac{d\mu}{d\zeta}\right)^2 = \frac{q^2 \gamma^4}{m^2 \kappa^2} \left(1 - [\kappa^2 m Z n_{\text{ion}} (v \sqrt{\mu^2 - \gamma^2} - \mu + \gamma) + 1]^{-2}\right) \quad (58)$$

and, using (58), consideration of the stationary points of μ yields $\gamma \leq \mu \leq \gamma^3(1+v^2)$. Furthermore, since $\mu(\zeta_{\text{I}}) = \gamma$ it follows $\frac{d\mu}{d\zeta} > 0$ for $\zeta_{\text{I}} < \zeta < \zeta_{\text{III}}$ where $\mu(\zeta_{\text{III}}) = \gamma^3(1+v^2)$ is the maximum value of μ (see figure 1). Thus, using (58) the period λ of the maximum amplitude oscillation is

$$\begin{aligned} \lambda &= 2(\zeta_{\text{III}} - \zeta_{\text{I}}) \\ &= \frac{2}{\omega_p \gamma^2} \int_{\gamma}^{\gamma^3(1+v^2)} \hat{\kappa} \frac{1}{\sqrt{1 - [\hat{\kappa}^2 (v \sqrt{\mu^2 - \gamma^2} - \mu + \gamma) + 1]^{-2}}} d\mu \end{aligned} \quad (59)$$

where $\hat{\kappa} = \kappa m \omega_p / |q|$.

A lower bound on λ may be determined by noting $0 \leq (v \sqrt{\mu^2 - \gamma^2} - \mu + \gamma) \leq \gamma - 1$ for $\gamma \leq \mu \leq \gamma^3(1+v^2)$. It follows

$$\frac{1}{\sqrt{1 - [\hat{\kappa}^2 (v \sqrt{\mu^2 - \gamma^2} - \mu + \gamma) + 1]^{-2}}} \geq \frac{1}{\sqrt{1 - [\hat{\kappa}^2 (\gamma - 1) + 1]^{-2}}} \quad (60)$$

and thus

$$\begin{aligned} \lambda &> \frac{2}{\omega_p \gamma^2} \int_{\gamma}^{\gamma^3(1+v^2)} \hat{\kappa} \frac{1}{\sqrt{1 - [\hat{\kappa}^2 (\gamma - 1) + 1]^{-2}}} d\mu \\ &= \frac{4m\gamma v^2}{|q| E_{\max}^{\text{BI}}}. \end{aligned} \quad (61)$$

Hence, using (57, 61) it follows λ diverges as least as fast as γ in the limit $v \rightarrow c$. Thus, although the Lorentz force on an electron trapped in a maximum amplitude wave is finite, the estimate $4m\gamma v^2/|q|$ of the work done on the electron over a wave period diverges in the limit $v \rightarrow c$ and, in principle, it is possible to accelerate electrons to arbitrarily high energies in the present classical theory. Furthermore, consideration of the electromagnetic stress-energy-momentum tensor $\mathcal{T}_{\text{EM}}$ arising from metric variations of the first term in (15) [3]

yields an electromagnetic mass-energy density ϱ_{EM} (in the lab frame) that diverges in the limit $v \rightarrow c$ for a wave on the verge of breaking. The tensor $\mathcal{T}_{\text{EM}}$ satisfies

$$\mathcal{T}_{\text{EM}}(T, U) = \frac{\partial \mathcal{L}_{\text{EM}}}{\partial X} \iota_T \star^{-1} \left(\iota_U F \wedge \star F - F \wedge \iota_U \star F \right) + \left(\mathcal{L}_{\text{EM}} - X \frac{\partial \mathcal{L}_{\text{EM}}}{\partial X} - Y \frac{\partial \mathcal{L}_{\text{EM}}}{\partial Y} \right) g(T, U) \quad (62)$$

where T, U are arbitrary vectors, and the 0-form ϱ_{EM} is

$$\varrho_{\text{EM}} = \mathcal{T}_{\text{EM}}(\partial/\partial x^0, \partial/\partial x^0). \quad (63)$$

Using (42, 62, 63) it follows

$$\varrho_{\text{EM}} = \left(2X \frac{\partial \mathcal{L}_{\text{EM}}}{\partial X} - \mathcal{L}_{\text{EM}} \right) \Big|_{Y=0} \quad (64)$$

and the Born-Infeld energy density ϱ_{BI} obtained using (64, 16),

$$\varrho_{\text{BI}} = \frac{1}{\kappa^2} \frac{1}{\sqrt{1 - \kappa^2 X}} (1 - \sqrt{1 - \kappa^2 X}), \quad (65)$$

diverges in the limit $E \rightarrow 1/\kappa$.

The behaviour of (59) for $\hat{\kappa}^2 \gamma \ll 1$ and $\gamma \gg 1$ may be extracted by changing the integration variable to $\chi = \mu/\gamma^3$. Using (59) it follows

$$\lambda = \frac{2\gamma}{\omega_p} \int_{\gamma^{-2}}^{1+v^2} \frac{1}{\sqrt{2I}} \left[1 + \frac{3}{4} \hat{\kappa}^2 I + \mathcal{O}(\hat{\kappa}^4 I^2) \right] d\chi \quad (66)$$

where

$$\begin{aligned} I &= \gamma^3 (v \sqrt{\chi^2 - \gamma^{-4}} - \chi + \gamma^{-2}) \\ &= \gamma \left(1 - \frac{1}{2} \chi \right) + \mathcal{O}(\gamma^{-1}). \end{aligned} \quad (67)$$

Since $\zeta = x^3 - vx^0$, the (angular) frequency ω_{BI} of the electrostatic oscillations measured in the lab frame is

$$\omega_{\text{BI}} = \frac{2\pi v}{\lambda} = \frac{2\pi}{\lambda} + \mathcal{O}(\gamma^{-2}) \quad (68)$$

and (66, 67, 68) yield

$$\omega_{\text{BI}} \approx \omega_{\text{AP}} \left[1 - \left(\frac{\kappa m \omega_p c}{2q} \right)^2 \gamma \right] \quad (69)$$

where the speed of light c has been restored and ω_{AP} is the (angular) frequency of electrostatic oscillations of a cold Maxwell plasma for $\gamma \gg 1$ [10],

$$\omega_{\text{AP}} = \frac{\pi}{2\sqrt{2}\gamma} \omega_p. \quad (70)$$

4 Conclusion

In practice, a typical laser-plasma experiment currently has $\gamma \sim 10-100$ and $\omega_p \sim 10^{14}$ rad/s; using $\kappa \sim 10^{-18}$ it follows $(\omega_{\text{BI}}/\omega_{\text{AP}} - 1) \sim 10^{-13} - 10^{-12}$. Although this value is currently out of reach of terrestrial experiments, we suggest that it may be possible to investigate the ramifications of (56, 61, 69) in the environment of astrophysical objects such as quasars and magnetars. In any case, a number of significant items have been omitted from the above calculation, such as the finite temperature of the plasma and ion motion. Future work will include more thorough evaluation of the consequences of non-zero κ in the context of more comprehensive plasma models.

Acknowledgements

We thank A Noble and RW Tucker for useful discussions, and thank the Cockcroft Institute of Accelerator Science and Technology for support. DAB is grateful for support provided by the ALPHA-X project. RMGMT is supported by the STFC Accelerator Science and Technology Centre (ASTEC).

References

- [1] GW Gibbons and CAR Herdeiro, Phys. Rev. D 63 (2001) 064006
- [2] R Ferraro, Phys. Rev. Lett. 99 (2007) 230401
- [3] T Dereli and RW Tucker, Euro. Phys. Lett. 89 (2010) 20009
- [4] M Born and L Infeld, Proc. R. Soc. Lond. A 144 (1934) 425
- [5] G Boillat, J. Math. Phys. 11 (1970) 941
- [6] J Plebanski, *Lectures on Non-Linear Electrodynamics*, Nordita, Copenhagen (1970)
- [7] ES Fradkin and AA Tseytlin, Phys. Lett. B 163 (1985) 123
- [8] HP Schlenvoigt, *et al.*, Nat. Phys. 4 (2008) 133
- [9] M Marklund, Nat. Photonics 4 (2010) 4
- [10] AI Akhiezer and RV Polovin, Sov. Phys. JETP 3 (1956) 696
- [11] JM Dawson, Phys. Rev. 113 (1959) 383
- [12] DA Burton and A Noble, J. Phys. A: Math. Theor. 43 (2010) 075502
- [13] RMGM Trines and PA Norreys, Phys. Plasmas 13 (2006) 123102
- [14] J Kijowski and G Magli, Class. Quantum Grav. 15 (1998) 3891
- [15] GA Maugin, *Material Inhomogeneities in Elasticity*, Applied Mathematics and Mathematical Computation, Chapman and Hall/CRC, (1993)
- [16] DA Burton, Theoret. Appl. Mech. 30 (2003) 85
- [17] <http://www.extreme-light-infrastructure.eu/>
- [18] <http://www.hiper-laser.org/>